

on galagos has (thus far) been observed only in a single immigrant matriline of the Kasakela community.

Red colobus, the prey of choice for many chimpanzee populations in East Africa, are not present at either Assirik or Fongoli where predation on galagos appears to be more common than elsewhere. McGrew⁶ hypothesized that chimpanzees at Mt. Assirik may face steep competition from other predators for most potential vertebrate prey (both primate and otherwise) within their range and so seek out arboreal, nocturnal prosimians as an alternative. The absence of red colobus at Fongoli may explain why galagos and other unusual prey such as mongoose¹², vervet monkeys¹³ and patas monkeys¹⁴ are consumed by chimpanzees there. The small size and nocturnal activity cycle of galagos may explain why they are consumed only rarely by chimpanzees of Kasakela and in the Mahale Mountains, given the availability of large, non-cryptic, diurnal prey such as red colobus.

ACKNOWLEDGEMENTS

I would like to thank Samson Pindu, Carson Murray, Anna Mosser, Matt Heintz and Emma Lantz for their friendship and assistance in the field, as well as Deus Mjungu, Anthony Collins and Shadrack Kamenya for advice and logistical support. Demographic data come from the long term records of JGI's Gombe Stream Research Centre. Leanne Nash and Thomas Butynski provided assistance with species identification. Craig Stanford, Anne Pusey, Ian Gilby and Jill Pruett provided helpful comments. I am grateful to the United Republic of Tanzania, particularly the Tanzania Commission for Science and Technology (COSTECH), the Tanzania Wildlife Research Institute (TAWIRI) and the Tanzania National Parks (TANAPA) for permission to conduct research at Gombe Stream NP. My field research was supported by a USC Joint Initiative Merit Fellowship Award and conducted under COSTECH permit 2009-229-ER-2007-188.

REFERENCES

- Stanford CB 1999. *The Hunting Apes: Meat-Eating and the Origins of Human Behavior*. Princeton University Press, Princeton.
- Goodall J 1986. *The Chimpanzees of Gombe: Patterns of Behavior*. Harvard University Press, Cambridge.
- Grubb P, Butynski TM, Oates JF, Bearder SK, Disotell TR, Groves CP, Struhsaker TT 2003. Assessment of the diversity of African primates. *Int J Primatol* **24**:1301–1357.
- Nishida T 1979. Predatory behavior among wild chimpanzees of the Mahale Mountains. *Primates* **20**:1–20.
- Uehara S 1997. Predation on mammals by the chimpanzee (*Pan troglodytes*). *Primates* **38**:193–214.
- McGrew WC 1983. Animal foods in the diets of wild chimpanzees: why cross-cultural variation? *J Ethol* **1**:46–61.
- Pruetz JD, Bertolani P 2007. Savanna chimpanzees (*Pan troglodytes verus*) hunt with tools. *Curr Biol* **17**:412–417.
- Butynski TM 1982. Blue monkey (*Cercopithecus mitis stuhlmanni*) predation on galagos. *Primates* **23**:563–566.
- Morris K, Goodall J 1977. Competition for meat between chimpanzees and baboons of the Gombe National Park. *Folia Primatol* **28**:109–121.
- Watts DP 2008. Scavenging by chimpanzees at Ngogo and the relevance of chimpanzee scavenging to early hominin behavioral ecology. *J Hum Evol* **54**:125–133.
- O'Malley RC, Wallauer B, Murray CM *in prep*. The appearance and spread of ant-fishing among Gombe chimpanzees: Inter-community transmission of a cultural behavior?

- Bogart SL, Pruett JD, Kante D 2008. Fongoli chimpanzee (*Pan troglodytes verus*) eats banded mongoose (*Mungos mungo*). *Pan Afr News* **15**:15–16.
- Gašperšič M, Pruett JD 2004. Predation on a monkey by savanna chimpanzees at Fongoli, Senegal. *Pan Afr News* **11**:8–10.
- Pruetz JD, Marshak JL 2009. Savanna chimpanzees (*Pan troglodytes verus*) prey on patas monkeys (*Erythrocebus patas*) at Fongoli, Senegal. *Pan Afr News* **16**:15–16.

* EDITORIAL NOTE:

PAN editors adopted the new trend in classification of the genus *Potamochoerus*, where there are two wild pig species: *P. porcus* (red river hog) and *P. larvatus* (bushpig), though the majority of researchers studying Eastern chimpanzees have followed a classic way of classifying bushpigs as *P. porcus*. Bushpigs (*P. larvatus*) are distributed mainly in East Africa, while red river hogs are distributed through the West and Central African rainforest belt (Wilson & Reeder 2005).

Wilson DE, Reeder DM (eds) 2005. *Mammal Species of the World: A Taxonomic and Geographic Reference (3rd ed)*. Johns Hopkins University Press, Baltimore.

<NOTE>

Chimpanzees at Semliki Ignore Oil Palms

William C. McGrew¹, Linda F. Marchant², Charlotte Payne¹, Timothy Webster³ & Kevin D. Hunt⁴

¹ Department of Biological Anthropology, University of Cambridge, UK

² Department of Anthropology, Miami University, USA

³ Department of Anthropology, Yale University, USA

⁴ Department of Anthropology, Indiana University, USA
(E-mail: wcm21@cam.ac.uk)

INTRODUCTION

Chimpanzee (*Pan troglodytes*) predation on oil palms (*Elaeis guineensis*) varies from essential to non-existent^{1,2}. At Gombe, in Tanzania, chimpanzees eat the pericarp of the fruit but discard the seed ('nut') containing the kernel; this foodstuff is the most frequently eaten item in their diet. At Mahale, also on the eastern shore of Lake Tanganyika, in Tanzania, the chimpanzees ignore all parts of the oil palm. However at Bossou, in Guinea, the apes consume the pericarp and pith, and use stone as hammer and anvil to crack open the dried fruit-shell, in order to consume the kernel. Bossou chimpanzees also show 'pestle-pounding', in which the detached petiole of a palm leaf is used to pulverize the apical meristem of an oil palm, before consumption.

The aim of this study was to see where on the spectrum of exploitation is the Semliki semi-habituated population of East African chimpanzees, as compared with Central³ and West African populations⁴⁻⁷. By regular monitoring and field experiment, we sought to test three hypotheses: 1) Oil palms are absent or unproductive at Semliki; 2) Raw materials for nut-cracking are absent or unsuitable at Semliki; 3) Oil palms and raw materials are

present and suitable, but the chimpanzees lack the cultural knowledge to make use of them.

METHODS

We studied the chimpanzees of Toro-Semliki Wildlife Reserve, Uganda⁸, recording both ecological and ethological data. We searched opportunistically and non-randomly for oil palms within the home range of the chimpanzees; searching was biased toward the vicinity of major trails used by both apes and researchers. Oil palms were not randomly dispersed, but instead occurred in groves (patches) of unknown origin; in this study we found such four groves. When we found an oil palm, it was described in terms of girth, height and morphology. Its location was noted by GPS and it was marked with plastic tape bearing basic information for identification. We cleared undergrowth and raked clean debris from an area of 5 m radius from the trunk, sufficient to be able to record fallen fruit and signs of predators visiting the tree, e.g. foot or knuckle prints.

Whenever we passed by, on an occasional, non-systematic basis, we monitored each palm's status, no



more than once daily. We noted its phenological status, e.g. fruiting, and any signs of predation, e.g. chewed fruits, stripped fronds, etc. Once data were taken, the area around the tree was swept clear again.

At each of the four groves, we assessed availability of potential hammers and anvils by running four 10-m × 1-m strip-transects from a central point in the grove in each of the cardinal directions (N, E, S, W). We recorded portable stones or wooden branches/sticks ('clubs') on the surface as potential hammers, and embedded stones, exposed tree roots, or fallen logs as potential anvils. For each stone, we recorded weight (to nearest 10 g), and length, width, and height (to nearest 1 cm). Stones weighing less than 200 g were ignored, as were small, unsuitable sticks or logs. We noted type of stone, when known, and stones were crudely classified as one of three shapes: sphere, pyramid, rectangular solid.

After a month of monitoring the palms, we devised an intervention in which we placed stones suitable to act as hammer and anvil at the base of the largest palm in each grove. Stones were chosen to match features reported from sites where chimpanzees crack nuts, such as Bossou. Nearby, we placed both fresh and dry nuts, and thereafter checked to see if the tools had been moved or the nuts cracked. Later, we placed nuts directly on the 'anvil' stone with hammer close by; that is, we tried to 'scaffold' the situation to encourage processing of the nuts.

RESULTS

Over the course of a 165-day field season, from 28 May to 8 Nov. 2008, we found 16 oil palm trees along the trail system in the Mugiri River valley. These ranged from a solitary tree to a grove of seven trees, making four groves in all (with the other three groves numbering three and four trees). To monitor the palms' status, we checked on them periodically, so that on average, each tree was visited 42 times (range: 12–69 visits), thus on average each tree was visited every 4 days (range: 2–14 days).

All palms had dried and de-husked nuts underneath their canopies. We did not try to count the total number of nuts, but most of them were empty shells, having been preyed upon by beetles (probably Bruchidae). When we collected and cracked intact nuts, they proved to be edible and indistinguishable from nuts tasted elsewhere.

Four of the palms bore fresh fruit during the study, in two of the four groves. They were the only palms of more than 10 m height (median: 13.5, range: 10–14 m) and girths of more than 100 cm (median: 117, range: 111–142). Their bright orange and yellow colouring made the fresh nuts conspicuous, and we counted each batch found on the ground, at each visit. One grove of four palms yielded fresh nuts in a season that lasted at least from 23 June to 16 July (on 15 visits); another grove of seven palms yielded nuts from 9 July to 18 Sept. (on nine visits). In the former case, the mean number of fresh nuts per visit was 17 (range: 1–61 fruits); the latter case the mean was 4 (range: 1–6 fruits). Found nuts were in one of three states: intact, chewed, or cracked/split open. So, who were the consumers? Several times we encountered monkeys and palm nut vultures (*Gypohierax angolensis*) in the palm canopy. We found no indirect signs of chimpanzees, but did find footprints of suids, probably bushpig (*Potamochoerus larvatus*).

None of the palms suffered feeding damage to any other part of the tree, including the fronds (leaves). This contrasts markedly with another palm species, *Phoenix reclinata*, the pith of which the Semliki chimpanzees ate daily, leaving tell-tale zigzag-shaped wadges strewn on the ground.

Potential raw materials were scarce on transects, compared with Lópe, Gabon³. Only 16 potential hammers and 7 potential anvils were found in the total of 160 m²

(4 × 4 × 10m²) of transected area, yielding an overall density of 0.1 hammers and 0.04 anvils per metre squared. Moreover, all 16 hammer stones were found in one 5-m stretch of eroding stream-bed. No other hammers or stones bigger than gravel, and no rocky outcrops were found anywhere else in the river valley, except in eroded gullies. The seven wooden anvils were either tree roots (5) or fallen logs (2). However, each grove had several hammer and anvil stones within 25–50 m of the palms, eroding out of gully sides and bottoms.

The 16 potential hammer stones found on transects averaged 317 g (range: 200–530) in weight, and had a mean length of 8 cm (range: 6–11), width of 6 cm (range: 4–7), and height of 4 cm (range: 3–6). A selection of eight stones found nearby in streambeds was bigger: mean weight: 822 g, length: 11 cm, width: 8.5 cm, height: 6 cm. Too few anvils were measured to yield findings on dimensions. Most stones in both sets were quartz, with which we easily cracked nuts. The most common shape of stone was rectangular solid.

DISCUSSION

We found no evidence that the chimpanzees of Semliki eat oil palms, in any form. So, why do they apparently ignore this valuable, potential food source? The presence of edible, productive and accessible oil palms, growing conveniently close to well-travelled chimpanzee trails provides strong evidence against hypothesis 1. Moreover, the Semliki chimpanzees *do* eat the fruits of *Phoenix reclinata* throughout their range, showing that they are not averse to Palmae fruits.

Although potential hammers and anvils were patchy in distribution, all groves had nearby sources of stones, at least, so that raw material scarcity cannot account for the absence of nut-cracking, at least by lithic elementary technology. Carriage of stones over distances of tens of metres, in order to crack oil palm nuts, is well known in West Africa⁹ and would have sufficed at Semliki. Thus, hypothesis 2 cannot account for the absence of nut-cracking.

Absence of oil palm exploitation seems unlikely to be environmentally precluded, thus leaving by exclusion support for hypothesis 3, that absence reflects cultural ignorance on the part of the apes^{2,10}. However, conclusions based on absence of evidence are always problematic, so more intensive and extensive study is needed.

ACKNOWLEDGEMENTS

Research permission was granted by the President's Office, Uganda National Council for Science and Technology, and Uganda Wildlife Authority. We thank the rangers and field assistants at Toro-Semliki Wildlife Reserve for their vital daily assistance.

REFERENCES

1. McGrew WC 1985. The chimpanzee and the oil palm: patterns of culture. *Soc Biol & Hum Affairs* **50**:7–23.
2. McGrew WC 1992. *Chimpanzee Material Culture: Implications for Human Evolution*. Cambridge University Press, Cambridge.
3. McGrew WC, Ham RM, White LTJ, Tutin CEG, Fernandez M 1997. Why don't chimpanzees in Gabon crack nuts? *Int*

J Primatol **18**:353–374.

4. Sugiyama Y, Koman J 1979. Tool using and making behavior in wild chimpanzees at Bossou, Guinea. *Primates* **20**:323–339.
5. Yamakoshi G, Sugiyama Y 1995. Pestle-pounding behavior of wild chimpanzees at Bossou, Guinea: a newly observed tool-using behavior. *Primates* **36**:489–500.
6. Biro D, Carvalho S, Matsuzawa T 2010. Tools, traditions, and technologies: interdisciplinary approaches to chimpanzee nut cracking. In: Lonsdorf EV, Ross SR, Matsuzawa T (eds), *The Mind of the Chimpanzee*. University of Chicago Press, Chicago, pp. 141–155.
7. Yamakoshi G 1998. Dietary responses to fruit scarcity of wild chimpanzees at Bossou, Guinea: Possible implications for ecological importance of tool use. *Am J Phys Anthropol* **106**:283–295.
8. Hunt KD, McGrew WC 2002. Chimpanzees in the dry areas of Assirik, Senegal, and Semliki Wildlife Reserve, Uganda. In: Boesch C, Hohmann G, Marchant LF (eds), *Behavioural Diversity in Chimpanzees and Bonobos*. Cambridge University Press, Cambridge, pp. 35–51.
9. Carvalho S, Cunha E, Sousa C, Matsuzawa T 2008. *Chaines operatoires* and resource exploitation strategies in chimpanzee nut-cracking (*Pan troglodytes*). *J Hum Evol* **55**:148–163.
10. McGrew WC 2004. *The Cultured Chimpanzee: Reflections on Cultural Primatology*. Cambridge University Press, Cambridge.

<NOTE>

A Case Report of Meat and Fruit Sharing in a Pair of Wild Bonobos

Satoshi Hirata¹, Shinya Yamamoto²,
Hiroyuki Takemoto² & Tetsuro
Matsuzawa²

1. Great Ape Research Institute, Hayashibara Biochemical Laboratories, Inc., Japan

2. Primate Research Institute, Kyoto University, Japan
(E-mail: hirata@gari.be.to)

INTRODUCTION

A number of observations of hunting and meat sharing in chimpanzees across different populations in Africa have been reported^{1,2}. In contrast, hunting is generally rare in bonobos (bilias). In Wamba, the Democratic Republic of the Congo, bonobos have been studied since 1973; however, only five cases of successful hunting have been reported, all of which involved a scaly-tailed squirrel as prey^{3,4,5}. Among these, two involved meat sharing. On the other hand, fruit sharing is fairly common among Wamba bonobos, and Kuroda argued that the sharing of highly preferred fruits among bonobos appears to resemble the meat sharing among chimpanzees⁶.

Fruit and meat sharing have also been documented in bonobos in Lomako⁷ (see refs. 8, 9, 10 for meat eating in other populations). When a case of sharing two species of fruit and a case of sharing a duiker were compared, the success rate of food transfer relative to the frequency of begging behavior was significantly lower in meat sharing.